

Resveratrol supplementation and acute pancreatitis: A comprehensive review

Shahram Agah^a, Abolfazl Akbari^{a,*}, Ehsan Sadeghi^b, Mojgan Morvaridzadeh^c,
Zarrin Basharat^d, Andriko Palmowski^e, Javad Heshmati^c

^a Colorectal Research Center, Iran University of Medical Sciences, Tehran, Iran

^b Research Center for Environmental Determinants of Health (RCEDH), Kermanshah University of Medical Sciences, Kermanshah, Iran

^c Department of Nutritional Science, School of Nutritional Science and Food Technology, Kermanshah University of Medical Sciences, Kermanshah, Iran

^d Jamil-ur-Rahman Center for Genome Research, Dr. Panjwani Center for Molecular Medicine and Drug Research, International Center for Chemical and Biological Sciences, University of Karachi, Karachi, 75270, Pakistan

^e Department of Rheumatology and Clinical Immunology, Charité – University Medicine Berlin, Germany

ARTICLE INFO

Keywords:

Resveratrol
Acute pancreatitis
Inflammation
Oxidative stress

ABSTRACT

Resveratrol, a natural polyphenolic ingredient extracted from herbs, suppresses oxidative stress and inflammation. We performed a comprehensive review to find any evidence about the effects of Resveratrol on acute pancreatitis (AP). Resveratrol has been found to directly impact cytokine generation. As these factors play a crucial role in the pathophysiology of AP, resveratrol might attenuate AP and its complications. Mechanistically, resveratrol exerts its pharmacological effects through anti-inflammatory and antioxidant mechanisms via interaction with different signaling molecules and transcription factors. Indeed, resveratrol might prove to be an effective therapeutic component for AP treatment in the future. In this review, we shed light on potential most recent pathways through which resveratrol might impact the management and control of AP.

1. Introduction

Acute pancreatitis (AP) is a severe inflammation of the pancreas causing enzyme activation inside the pancreas, triggering auto-degradation accompanied by local and systemic inflammation [1]. The prevalence of AP is increasing worldwide [2]. Despite advances in diagnostics and treatment, AP is still associated with both severe morbidity and mortality [3]. Multiple factors may be involved in the pathogenesis of AP, e.g., alcohol consumption or bile duct obstruction. If causal factors are taken care off immediately, both the pancreatic structure and activity can recover to normal. Still, AP may become a life-threatening condition [4]. For instance, severe AP can alter the function and structure of the pancreatic duct and eventually cause chronic obstructive pancreatitis [5]. Current state-of-the-art care of AP mainly consists of supportive care including monitoring vital signs and arterial oxygen saturation, hydration, and analgesic treatment [6].

Resveratrol (3,5,40-trihydroxy-transstilbene), a well-known traditional medicine, it is a polyphenol rich ingredient extracted from several plants [7]. The highest concentration of resveratrol (50–100 µg/g) is found in grape skins [8,9]. Resveratrol is usually found in herbs in

trans-resveratrol form, and when orally consumed, trans-resveratrol is quickly transformed to the biologically more active form of dihydro-resveratrol (Fig. 1) [10]. Resveratrol has several biological properties, e.g., it inhibits platelet aggregation [11] and exerts anti-inflammatory, chemopreventive and antioxidant effects [12]. Because of its wide range of pharmacological effects and easy methods of extraction, resveratrol has been used in the management of a lot of diseases and received an increasing amount of attention in recent years [13]. It has been shown in previous studies that supplementation of resveratrol has an effect on proinflammatory cytokines, hyperglycemia, and antioxidant competence and protects β-cell ultrastructure [14]. Benefits of resveratrol in the management of AP have been suggested because of its antioxidant, anti-inflammatory, and immunomodulatory functions [15]. Resveratrol exerts its effects through modulation of several metabolic factors, transcriptional elements, and miRNAs (Fig. 2).

AP is often accompanied by a systemic inflammatory response syndrome due to an elevation of inflammatory mediators requiring urgent management [16,17]. Resveratrol has antioxidative and anti-coagulative and immunomodulatory properties, all of which may alleviate AP. The aim of this review was to summarize the most recent

* Corresponding author at: Colorectal Research Center, Rasoul-e Akram Hospital, Sattarkhan Ave, Niyayesh St., Tehran, Iran.

E-mail addresses: akbari.ab@iums.ac.ir (A. Akbari), Javad.heshmati@gmail.com (J. Heshmati).

<https://doi.org/10.1016/j.bioph.2021.111268>

Received 29 September 2020; Received in revised form 7 January 2021; Accepted 10 January 2021

Available online 22 January 2021

0753-3322/© 2021 Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. Resveratrol and its bioconversion.




Fig. 2. Resveratrol and the factors it may possibly interact with in acute pancreatitis. Cyclooxygenase-2 (COX-2); Phosphoinositide 3-kinases (PI3Ks); Inducible nitric oxide synthase (iNOS); Endothelial nitric oxide synthase (eNOS); Nitric oxide (NO); MicroRNAs (miRNAs); Myeloperoxidase (MPO); Heat shock factor-1 (HSF-1); Sirtuin 1 (SIRT1); Superoxide dismutase (SOD); Glutathione peroxidase (GSHPx); Reactive oxygen species (ROS); B-cell lymphoma 2 (Bcl-2); Intercellular Adhesion Molecule (ICAM); vascular cell adhesion molecule (VCAM); Interleukins (ILs); Endothelin (ET); Tumor necrosis factor alpha (TNF α); 2,4-Dinitrophenylhydrazine (DNPH); nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B); Malondialdehyde (MDA).

findings about the effects of resveratrol on AP.

1.1. Antioxidant effects of resveratrol

Histopathological results showed that treatment with resveratrol preserves the pancreatic acinar cells from degeneration, leukocyte infiltration and edema in rats [18–20]. There is growing evidence indicating the pivotal role of free radicals in the pathophysiology of several inflammatory conditions, including AP [21]. It has been shown that oxidative stress is one main reason for acinar cell damage in experimental AP [22]. Moreover, it seems that ROS cause increasing pain in patients suffering from AP [23]. Recent findings indicate that AP causes a decrease of strength in the antioxidant defense system, for instance, it has been shown that total antioxidant status (TAS) is decreased in AP patients. Interestingly, research has shown that resveratrol can restore the TAS to normal levels in AP patients [24]. It has been reported that oral administration of resveratrol significantly inhibits glutathione depletion in cerulein-induced AP rats [25,26].

As AP causes an oxidative alteration of proteins, elevation of the amount of 2,4-dinitrophenylhydrazine (DNPH)-reactive protein carbonyls was observed in the pancreatic tissue of AP rats [27,28]. Interventions with resveratrol were shown to significantly diminish irreversible carbonylation of proteins during AP [29,30]. In addition, some studies showed that resveratrol restored Plasma glutathione peroxidase (GSH-Px) concentrations and reduced lipid peroxidation in the pancreas [20,31,32]. Lipid peroxidation leads to increased damage in lipid membranes, such as those of zymogen granules in the pancreas [33,34]. Malondialdehyde (MDA) is the degraded organic compound of lipid peroxidation which is generated when unsaturated fatty acids of cell membranes are attacked by ROS, consequently, MDA concentrations

can reliably and indirectly indicate the level of lipid peroxidation in tissues [35]. There are several studies which indicate that resveratrol significantly reduced MDA levels in AP patients, proposing that resveratrol efficiently suppresses lipid peroxidation [25,36]. Resveratrol consumption has protective effects on the gastrointestinal mucosa, possibly due to its antioxidant activities and radical scavenging. This is probably a major mechanism of resveratrol against gastrointestinal injuries caused by AP. Superoxide dismutase (SOD) is another antioxidant enzyme which can decrease oxidative stress and inflammatory activity [37]. Resveratrol has been shown to increase SOD levels in several studies [38,39]. In addition, resveratrol has effects on other mediators of oxidative stress such as Nitric oxide (NO) and Myeloperoxidase (MPO). NO is present in several physiological processes such as neurotransmission, inflammation and modulation of vascular tone. For instance, resveratrol has been shown to have a cardioprotective effect by elevating the generation of endothelial Nitric oxide synthase (eNOS) [40], inducible nitric oxide synthase (iNOS) [41] and NO [42]. However, other studies found that resveratrol decreased the expression of iNOS and NO [43,44]. Although the true function of NO in the pathophysiology of AP has not been totally understood, eNOS has been demonstrated to have a protective impact by elevating the pancreatic blood flow rate [45]. At the same time, however, iNOS also shows pro-inflammatory functions [46], and it has been shown that serum concentrations of NO are associated with increasing severity and mortality from AP in randomized clinical trials [47].

The production of superoxide anion, catalysed by NADPH oxidase, is increased as one of the main outcomes of oxidative stress [48]. It has been demonstrated that the NADPH oxidase system activity increased significantly in AP [49,50]. Oral consumption of dihydro-resveratrol has been shown to decrease, in a considerable amount, the NADPH oxidase

regulates proinflammatory cytokines [54]. Tumor necrosis factor alpha (TNF- α), which is secreted from polymorphonuclear neutrophils, lymphocytes and macrophages [55], is considered to be a “first-line” cytokine [56]. TNF- α induces the generation of other pro-inflammatory factors and oxidant factors including ROS, and it also stimulates the oxidative stress-responsive genes that extend the chronic inflammatory response [57]. Furthermore, TNF- α synergistically impacts the generation of nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) and other cytokines related to inflammation [58]. NF- κ B activated genes can induce cascade sequence-associated genes such as Interleukin 6 (IL-6), TNF- α , Interleukin-8 (IL-8), Interleukin-2 (IL-2), Intercellular Adhesion Molecule 1 (ICAM-1), etc [59]. Adhesion molecules such as ICAM-1 and vascular cell adhesion molecule 1 (VCAM-1) have a critical role in the activation of inflammatory response and leukocyte adherence [60]. Their gene expression is stimulated via induction of inflammatory mediators, including TNF- α and IL-1 [61]. TNF- α modulates the gene expression of VCAM-1 and ICAM-1 in the gastrointestinal tract and regulates the adherence of monocytes, neutrophils, and lymphocytes to vascular endothelial cells. These cells, especially neutrophils, may cause damage to gastrointestinal endothelial cells through degranulation [62]. Recent findings show that administration of resveratrol alleviates injuries to the intestinal mucosal barrier by inhibiting ICAM-1 and VCAM-1 surface gene expression [63,64]. In addition, it has been demonstrated that resveratrol is able to suppress the activation of the activator protein-1 (AP-1) [24] and NF- κ B [65]. AP-1 is a major transcription element which modulates various cytological functions, such as oncogenic transformation, apoptosis, differentiation, cell death, cell migration, and proliferation along with progression in various tissues [66]. Correspondingly, resveratrol has been shown to be related with a reduced level of IkappaB alpha (IkB α) degradation and suppression of AP-1 functions [67]. It has been shown that this suppression of NF- κ B via resveratrol may be responsible for the inhibition of cell-mediated cytotoxicity and lymphocyte proliferation [68]. In the cytosol, inhibitor protein kappa B (IkB) binds to NF- κ B and deactivates it. After activation of IkB, specific IkB kinases phosphorylate IkB; afterwards, IkB is rapidly degraded through proteasome-dependent pathways [69]. It has been shown that resveratrol inhibits the activation of NF- κ B by suppressing the degradation of IkB [70].

Furthermore, it has been indicated that increased positive signals of Phosphoinositide 3-kinases (PI3K p85) were reported in pancreatic tissues, and the intensity of PI3K signals decreased after resveratrol administration [71]. Moreover, Akt (Protein Kinase B) phosphorylation was observed to be increased in AP tissues. It also diminished after resveratrol administration [25,72]. Besides, antioxidant factors including resveratrol have the capability to decrease the level of leukocyte transfer to damaged tissue by intervening with myeloperoxidase function [73]. It seems that the immunomodulatory characteristics of resveratrol, could be more effective in amelioration of AP than inhibition of inflammatory factors.

Resveratrol administration does not only decrease inflammatory mediators but also increases anti-inflammatory cytokines. For example, some studies reported that resveratrol intervention upregulated the expression of IL-10 [74]. IL-10 is known to have anti-inflammatory properties due to its effect on the generation of pro-inflammatory cytokines [75]. The effect of resveratrol on inflammatory factors is presented in Fig. 4.

1.3. Effects of resveratrol on tissue damage due to AP

Formation of pancreatic edema is one of the main histological criteria for pancreatic damage [76]. It has been shown that resveratrol administration significantly reduces pancreatic edema in experimental models [25]. Severe, end stage forms of AP are determined by the progression of parenchymal necrosis accompanied by systemic complications and leukocyte infiltration, all of which are related to a worsening of the clinical condition of the patient [77]. It has been shown that

resveratrol reduces leukocyte infiltration, but it does not significantly alter tissue damage [18,78].

Severe forms of AP usually cause several systemic complications which even affect distant organs including the lungs, heart, brain, liver, kidneys, and bones [79]. Acute lung damage is regarded as one of the most recurrent potentially destructive complications of severe AP [80]. However, the pathogenesis of AP-related lung damage is still not totally understood, but it is probably related to an increase of pro-inflammatory cytokines in the lung. Recent findings indicate that oral consumption of resveratrol significantly improves AP-related general inflammatory reactions in the lungs through inhibition of the NF- κ B signaling pathway and leads to a significant decrease of pro-inflammatory mediators [81]. Actually, pulmonary wall thickening, causing alveolar obstruction, is the most frequent histological feature of AP-associated systemic inflammatory damage [82]. Resveratrol administration, even at low doses such as 10 mg/kg body weight, is associated with improvement of pulmonary wall thickening [81,83]. Amelioration of histological lung damage scores by administration of resveratrol is mainly related to inhibitory effects of resveratrol on the generation of different pro-inflammatory mediators such as, IL-6, IL-1 β and TNF- α [84].

Heart failure is another cause of death in AP patients [85]. Microcirculation disturbance plays a crucial role in chronic heart complications [86]. Previous investigations have shown that microcirculatory disturbance also has a pivotal role in the pathogenesis of AP [87]. Based on recent findings, it can be proposed that resveratrol administration reduces microcirculatory disturbance and suppresses necrotic cell death during AP [88]. Experimental studies suggest several mechanisms for the effect of resveratrol on microcirculatory disturbance, including a decrease of edema and inflammation via down-regulation of Apoptosis regulator BAX (Bax) and caspase 3 and upregulation of Bcl-2 [89]. Moreover, resveratrol diminishes gene expression of cytochrome C and keeps cells safe from injuries [90,91].

Previous studies report that plasma renin activity and Angiotensin II (Ang II) concentrations rise during the progression of AP. It has been demonstrated that the renin-angiotensin system has a crucial role in the initiation of AP [26], and Ang II, plasma endothelin (ET), and NO are up-regulated during AP [92]. In addition to beneficial role of resveratrol on microcirculatory disturbance, investigators show that resveratrol administration improves plasma renin activity and down-regulates plasma ET, Ang II, and NO [88]. Researchers propose that plasma levels of ET and NO are increased in initial stages of AP and change the amount of blood flow in local pancreatic vessels. Altogether, these alterations cause hypoxia, hemodynamic instability, tissue ischemia, and finally, advanced AP [93]. Elevated concentration of ET upregulate intracellular calcium and increase the function of pro-inflammatory cells [94]. Plasma concentration of NO is also elevated because of increased NOS-activity [95]. Resveratrol administration is likely to diminish concentrations of ET and NO via suppression of ET receptors and NOS [96]. Previous investigations proposed that resveratrol suppresses the expression of iNOS and decreases inflammation and pancreatic tissue injury [97].

It has been shown that, Resveratrol also can improve the efficacy and therapeutic effects of other treatment approaches for AP. For example Bone marrow-derived mesenchymal stem cells (BMSCs) are recently proposed to treatment of AP [98]. It has been demonstrated that resveratrol pretreatment of BMSCs (Res-BMSCs) improve the therapeutic impact of decreasing apoptosis of pancreatic cells and increase regeneration of damaged blood vessels [71].

1.4. Resveratrol and enzymes in AP

Hyperlipidemia and hyperamylasemia are typical features of initial early-stage AP [99]. Indeed, these features are often used for the diagnosis of AP [100]. Recent findings shed light on the role of resveratrol supplementation concerning premature activation of pancreatic enzymes [52]. Increased α -amylase is a primary diagnostic criterion for AP




Fig. 5. Resveratrol decreases damage to pancreatic tissue through suppression of calcium overload; it suppresses calcium overload and thereby decreases trypsinogen activation, oxidative stress, mitochondrial dysfunction and microcirculatory disorders; Resveratrol also decreases damage to other organs including lung and heart by decreasing microcirculatory dysfunction.

[101]. Although amylase and lipase enzymes often increase threefold in AP patients, there seems to be no association between the severity of AP and serum levels of these enzymes [102]. However, it has been reported that resveratrol administration can reduce levels of serum amylase and lipase in AP patients [20,25,29]. Moreover, recent experimental studies indicate that intervention with resveratrol was not enough to suppress tissue damage, although amylase and lipase serum concentrations decreased [67]. As serum amylase concentration is the main biomarker for AP, serum Lactate dehydrogenase (LDH) activity could represent the amount of necrosis [103] and MPO function is associated with neutrophil recruitment [104]. Serum pancreatic MPO and LDH activity are remarkably elevated in AP. Resveratrol diminishes pancreatic MPO activity and serum LDH activity in rat AP-models [105].

Trypsin activation due to pro-inflammatory cytokine-release is recognized to be another initial mechanism in AP-mediated tissue damage [106,107]. However, experimental studies show that resveratrol does not suppress trypsin activity [67].

1.5. Resveratrol and gene expression in AP

mRNA gene expression of inflammatory mediators such as IL-6, TNF- α , IL-10 is thought to change in AP patients [108]. Resveratrol was found not only to significantly diminish pancreatic IL-6 and TNF- α mRNA expression, but it also increases pancreatic IL-10 mRNA expression [105]. Sirtuin 1 (SIRT1) is a NAD dependent deacetylase that modulates cell proliferation, metabolism, and DNA repair and acts as a modulator for several cellular antioxidant processes [109]. Recent findings show that SIRT1 modulates the expression of specific glucose sensing and insulin secretion in the pancreas through mitochondria-related genes that control metabolic coupling [110]. Resveratrol was found to be a classic SIRT1 stimulator [111], and it has been shown that resveratrol protects pancreatic cells by increased expression of SIRT1 [112,113]. Overexpression of SIRT1 gene seems to protect pancreatic β -cells from toxicity from inflammatory mediators by inhibiting the NF- κ B signaling pathway [114]. It has been shown that resveratrol also protects

pancreatic β -cells by augmenting SIRT1 [115].

P53 is a pro-apoptotic protein which is strongly associated with SIRT1. p53 expression was remarkably decreased in the pancreas of rats with AP [116], and resveratrol increased p53 expression significantly [117]. Moreover, HSF1, which is one of the main heat shock transcription elements, is expressed widely in different kind of cells and has a pivotal role in inflammation by modulating the transcription of relevant cytokines and inflammatory mediators [118–120]. Acetylation of HSF1 diminishes its DNA attachment function and thereby increases transcription of pro-inflammatory mediators and suppresses transcription of anti-inflammatory elements. It has been shown that HSF1-knockout mice with pancreatitis induced by cerulein experienced acute edema that was more severe than in control mice [83]. It has been demonstrated that HSF1 expression was remarkably reduced in pancreatic tissues of AP mice, indicating that HSF1 has a role in the pathogenesis of AP. Resveratrol does not only increase the total expression of HSF1, but resveratrol administration also decreases acetylated HSF1 [105].

In addition, resveratrol exerts its protective effects against AP through microRNA (miRNA) modification. It seems that resveratrol exerts its antitumor and anti-inflammatory effects through modulation of the expression of specific miRNAs [121]. For instance, the expression level of miR-126 was upregulated by administration of resveratrol [122]. miR-126 suppresses inflammation and inflammatory mediators in endothelial cells [123]. Resveratrol changes miR-126 expression through upregulation of KLF2. KLF2 is a zinc-finger family's transcription factor; a crucial regulator in the progress of cell development and differentiation [124]. miR-21 is a necessary element for ROS-promoted pancreatic stellate cell function, and decrease in miR-21 interrupts ROS-promoted migration in pancreatic stellate cells [125]. Resveratrol was found to suppress pancreatic cancer cell migration and invasion via inhibition of glycolysis in pancreatic stellate cells and ROS/miR-21-mediated function [126]. Resveratrol also diminishes the expression of miR-21 and inhibits BCL-2 expression and thereby protects pancreatic cells from necrosis [127].

1.6. Effects of resveratrol on calcium overload

Several triggers have been recognized to be involved in the etiology of AP, but exact mechanisms are still poorly understood. Free calcium ions inside the cytoplasm of acinar cells function as a main second messenger. Calcium overload may have pivotal role in the pathogenesis of AP [128]. Calcium signaling pathways have been shown to be disturbed in initial stages of AP [129]. Under normal and healthy circumstances, calcium concentrations are stable, while in pathological situations, several factors can impact intracellular calcium concentrations and may lead to a rise in intracellular calcium levels causing a calcium overload [130]. This may be true for AP as well [131]. Recent research shows that calcium overload is not only an etiological indicator for AP, it also worsens the condition by effects on other organs and organelles [129]. It has been shown that cytoplasmic calcium overload causes dysfunction of acinar cell secretion, increased activation of trypsinogen, early activation of zymogens, cell necrosis, oxidative stress, microcirculatory dysfunctions, and mitochondrial disorders [128]. Intracellular calcium overload is also associated with severe lung tissue damage in AP patients. Resveratrol administration can efficiently decrease the complications of AP by regulating intracellular calcium-modulatory pathways and by decreasing calcium overload [132]. The effect of resveratrol on calcium overload and tissue damage is presented in Fig. 5.

2. Conclusions

Acute pancreatitis is a common, severe and potentially life-threatening acute gastrointestinal disease. Recent evidence suggests that resveratrol might be a useful tool in the management of AP. Resveratrol might alleviate pancreatic damage and damage in other organs through various pathways. It seems to suppress NF- κ B activity and regulate the generation of inflammatory mediators, and it improves the anti-oxidative defense system, modulates and manages intracellular calcium in pancreatic acinar cells and modulates the expression of several genes and transcriptional factors that are related to the pathophysiology of AP. To sum up, resveratrol seems to have several beneficial effects on AP, but high-quality trials in humans are needed to underpin these findings.

Credit author statement

A.A and contributed to the Conceptualization, Investigation, Methodology, Validation and Visualization. S.H.A. contributed to Project administration, Supervision, Validation and Visualization. J.H. and A.P contributed to Writing - original draft. M.M and Z.B. contributed to Writing - review & editing.

Funding information

The authors receive no funding for this research

Transparency document

The [Transparency document](#) associated with this article can be found in the online version.

Declaration of Competing Interest

The authors declare no conflict of interest.

Acknowledgements

This work was supported by Grant No. 97-4-49-13925 from Iran University of Medical Sciences. We also thank Alireza Mirzaei cooperation, Iran University of medical sciences, Tehran, Iran for helping us

with the manuscript revision.

References

- [1] S.M. Van Dijk, N.D. Hallensleben, H.C. van Santvoort, P. Fockens, H. van Goor, M.J. Bruno, M.G. Besselink, Acute pancreatitis: recent advances through randomised trials, *Gut* 66 (11) (2017) 2024–2032.
- [2] Q. Wang, M. Shen, X. Leng, X. Zeng, F. Zhang, J. Qian, Prevalence, severity, and clinical features of acute and chronic pancreatitis in patients with systemic lupus erythematosus, *Rheumatol. Int.* 36 (10) (2016) 1413–1419.
- [3] S.J. Hashemi, R. Masjedizadeh, F. Mehran, A. Tahmasebi, The relationship between etiology of acute pancreatitis and its clinical features in Iranian patients, *Biomed. Pharmacol. J.* 14 (2) (2017) 875–880.
- [4] M. Borran, M. Minaian, B. Zolfaghari, P. Mahzouni, Protective effect of Tribulus terrestris fruit extract on cerulein-induced acute pancreatitis in mice, *Avicenna J. Phytomed.* 7 (3) (2017) 250.
- [5] A.P. Shah, M.M. Mourad, S.R. Bramhall, Acute pancreatitis: current perspectives on diagnosis and management, *J. Inflamm. Res.* 11 (2018) 77.
- [6] S.D. Crockett, S. Wani, T.B. Gardner, Y. Falck-Ytter, A.N. Barkun, S. Crockett, J. Feuerstein, S. Flamm, Z. Gellad, L. Gerson, American Gastroenterological Association Institute guideline on initial management of acute pancreatitis, *Gastroenterology* 154 (4) (2018) 1096–1101.
- [7] S. Abdollahi, A. Salehi-Abargouei, O. Toupchian, M.H. Sheikhha, H. Fallahzadeh, M. Rahmani, M. Tabatabaie, H. Mozaffari-Khosravi, The effect of resveratrol supplementation on cardio-metabolic risk factors in patients with type 2 diabetes: a randomized, double-blind controlled trial, *Phytother. Res.* 33 (12) (2019) 3153–3162.
- [8] E. Wenzel, V. Somoza, Metabolism and bioavailability of trans-resveratrol, *Mol. Nutr. Food Res.* 49 (5) (2005) 472–481.
- [9] U. Gasić, I. Čirić, T. Pejčić, D. Radenković, V. Djordjević, S. Radulović, Ž. Tesić, Polyphenols as possible agents for pancreatic diseases, *Antioxidants* 9 (6) (2020) 547.
- [10] L.D. Williams, G.A. Burdock, J.A. Edwards, M. Beck, J. Bausch, Safety studies conducted on high-purity trans-resveratrol in experimental animals, *Food Chem. Toxicol.* 47 (9) (2009) 2170–2182.
- [11] C. Bonechi, S. Lamponi, A. Donati, G. Tamasi, M. Consumi, G. Leone, C. Rossi, A. Magnani, Effect of resveratrol on platelet aggregation by fibrinogen protection, *Biophys. Chem.* 222 (2017) 41–48.
- [12] C. Buhmann, P. Shayan, B. Popper, A. Goel, M. Shakibaei, Sirt1 is required for resveratrol-mediated chemopreventive effects in colorectal cancer cells, *Nutrients* 8 (3) (2016) 145.
- [13] A.Y. Berman, R.A. Motechin, M.Y. Wiesenfeld, M.K. Holz, The therapeutic potential of resveratrol: a review of clinical trials, *NPJ Precis. Oncol.* 1 (1) (2017) 1–9.
- [14] P. Palsamy, S. Subramanian, Ameliorative potential of resveratrol on proinflammatory cytokines, hyperglycemia mediated oxidative stress, and pancreatic β -cell dysfunction in streptozotocin-nicotinamide-induced diabetic rats, *J. Cell. Physiol.* 224 (2) (2010) 423–432.
- [15] K. Farhadi, F. Esmailzadeh, M. Hatami, M. Forough, R. Molaie, Determination of phenolic compounds content and antioxidant activity in skin, pulp, seed, cane and leaf of five native grape cultivars in West Azerbaijan province, Iran, *Food Chem.* 199 (2016) 847–855.
- [16] X. Chen, B. Ji, B. Han, S.A. Ernst, D. Simeone, C.D. Logsdon, NF- κ B activation in pancreas induces pancreatic and systemic inflammatory response, *Gastroenterology* 122 (2) (2002) 448–457.
- [17] B.J. John, S. Sambandam, P. Garg, G. Singh, M. Kaur, R. Baskaran, G. Srinivasan, V. Leelakrishnan, K. Janarthan, Persistent systemic inflammatory response syndrome predicts the need for tertiary care in acute pancreatitis, *Acta Gastroenterol. Belg.* 80 (3) (2017) 377–380.
- [18] C. Carrasco, M.S. Holguín-Arévalo, G. Martín-Partido, A.B. Rodríguez, J. A. Pariente, Chemopreventive effects of resveratrol in a rat model of cerulein-induced acute pancreatitis, *Mol. Cell. Biochem.* 387 (1–2) (2014) 217–225.
- [19] E. Gedik, S. Girgin, H. Ozturk, B.D. Obay, H. Ozturk, H. Buyukbayram, Resveratrol attenuates oxidative stress and histological alterations induced by liver ischemia/reperfusion in rats, *World J. Gastroenterol. WJG* 14 (46) (2008) 7101.
- [20] Z.-D. Li, Q.-Y. Ma, C.-A. Wang, Effect of resveratrol on pancreatic oxygen free radicals in rats with severe acute pancreatitis, *World J. Gastroenterol. WJG* 12 (1) (2006) 137.
- [21] S. Bopanna, B. Nayak, S. Prakash, S.J. Mahapatra, P.K. Garg, Increased oxidative stress and deficient antioxidant levels may be involved in the pathogenesis of idiopathic recurrent acute pancreatitis, *Pancreatol.* 17 (4) (2017) 529–533.
- [22] E. Özkan, C. Akyüz, E. Dulundu, Ü. Topaloglu, A.O. Şehirli, F. Ercan, G. Şener, Protective effects of lycopene on cerulein-induced experimental acute pancreatitis in rats, *J. Surg. Res.* 176 (1) (2012) 232–238.
- [23] Y. Irie, M. Tsubota, H. Ishikura, F. Sekiguchi, Y. Terada, T. Tsujiuchi, K. Liu, M. Nishibori, A. Kawabata, Macrophage-derived HMGB1 as a pain mediator in the early stage of acute pancreatitis in mice: targeting RAGE and CXCL12/CXCR4 axis, *J. Neuroimmune Pharmacol.* 12 (4) (2017) 693–707.
- [24] S.K. Manna, A. Mukhopadhyay, B.B. Aggarwal, Resveratrol suppresses TNF-induced activation of nuclear transcription factors NF- κ B, activator protein-1, and apoptosis: potential role of reactive oxygen intermediates and lipid peroxidation, *J. Immunol.* 164 (12) (2000) 6509–6519.

- [25] S.W. Tsang, Y.-F. Guan, J. Wang, Z.-X. Bian, H.-J. Zhang, Inhibition of pancreatic oxidative damage by stilbene derivative dihydro-resveratrol: implication for treatment of acute pancreatitis, *Sci. Rep.* 6 (2016) 22859.
- [26] A. Kode, S. Rajendrasozhan, S. Caito, S.-R. Yang, I.L. Megson, I. Rahman, Resveratrol induces glutathione synthesis by activation of Nrf2 and protects against cigarette smoke-mediated oxidative stress in human lung epithelial cells, *Am. J. Physiol.-Lung Cell. Mol. Physiol.* 294 (3) (2008) L478–L488.
- [27] T. Reinheckel, B. Nedelev, J. Prause, W. Augustin, H.-U. Schulz, H. Lippert, W. Halangk, Occurrence of oxidatively modified proteins: an early event in experimental acute pancreatitis, *Free Radic. Biol. Med.* 24 (3) (1998) 393–400.
- [28] C.C. Winterbourn, M.J. Bonham, H. Buss, F.M. Abu-Zidan, J.A. Windsor, Elevated protein carbonyls as plasma markers of oxidative stress in acute pancreatitis, *Pancreatology* 3 (5) (2003) 375–382.
- [29] M. Lawinski, Z. Sledzinski, J. Kubasik-Juraniec, J.H. Spodnik, M. Wozniak, W. Boguslawski, Does resveratrol prevent free radical-induced acute pancreatitis? *Pancreas* 31 (1) (2005) 43–47.
- [30] P. Palsamy, S. Sivakumar, S. Subramanian, Resveratrol attenuates hyperglycemia-mediated oxidative stress, proinflammatory cytokines and protects hepatocytes ultrastructure in streptozotocin–nicotinamide-induced experimental diabetic rats, *Chem. Biol. Interact.* 186 (2) (2010) 200–210.
- [31] Y.-E. Lee, J.-W. Kim, E.-M. Lee, Y.-B. Ahn, K.-H. Song, K.-H. Yoon, H.-W. Kim, C.-W. Park, G. Li, Z. Liu, Chronic resveratrol treatment protects pancreatic islets against oxidative stress in db/db mice, *PLoS One* 7 (11) (2012).
- [32] A. Baltaci, H. Gokbudak, S. Baltaci, R. Mogulkoc, M. Avunduk, The effects of resveratrol administration on lipid oxidation in experimental renal ischemia-reperfusion injury in rats, *Biotech. Histochem.* 94 (8) (2019) 592–599.
- [33] C. Col, K. Dinler, O. Hasdemir, O. Buyukasik, G. Bugdayci, Oxidative stress and lipid peroxidation products: effect of pinealectomy or exogenous melatonin injections on biomarkers of tissue damage during acute pancreatitis, *Hepatobiliary Pancreat. Dis. Int.* 9 (1) (2010) 78–82.
- [34] C. Niederau, H. Klonowski, H.-U. Schulz, M. Sarbia, R. Lüthen, D. Häussinger, Oxidative injury to isolated rat pancreatic acinar cells vs. isolated zymogen granules, *Free Radic. Biol. Med.* 20 (7) (1996) 877–886.
- [35] G. Barrera, S. Pizzimenti, M. Daga, C. Dianzani, A. Arcaro, G.P. Cetrangolo, G. Giordano, M.A. Cucci, M. Graf, F. Gentile, Lipid peroxidation-derived aldehydes, 4-hydroxynonenal and malondialdehyde in aging-related disorders, *Antioxidants* 7 (8) (2018) 102.
- [36] R.K. Jha, Q. Ma, Z. Lei, H. Sha, Resveratrol ameliorates the deleterious effect of severe acute pancreatitis, *Cell Biochem. Biophys.* 62 (2) (2012) 397–402.
- [37] P. Kangari, T.Z. Farahany, A. Golchin, S. Ebadollahzadeh, A. Salmani, A. Mahboob, A. Nourazarian, Enzymatic antioxidant and lipid peroxidation evaluation in the newly diagnosed breast cancer patients in Iran, *Asian Pacific J. Cancer Prev.: APJCP* 19 (12) (2018) 3511.
- [38] W. Li, L. Cao, X. Chen, J. Lei, Q. Ma, Resveratrol inhibits hypoxia-driven ROS-induced invasive and migratory ability of pancreatic cancer cells via suppression of the Hedgehog signaling pathway, *Oncol. Rep.* 35 (3) (2016) 1718–1726.
- [39] T. Szkudelski, K. Szkudelska, Potential of resveratrol in mitigating metabolic disturbances induced by ethanol, *Biomed. Pharmacother.* 101 (2018) 579–584.
- [40] T. Wallerath, G. Deckert, T. Ternes, H. Anderson, H. Li, K. Witte, U. Förstermann, Resveratrol, a polyphenolic phytoalexin present in red wine, enhances expression and activity of endothelial nitric oxide synthase, *Circulation* 106 (13) (2002) 1652–1658.
- [41] F. Pellegatta, A.A. Bertelli, B. Staels, C. Duhem, A. Fulgenzi, M.E. Ferrero, Different short- and long-term effects of resveratrol on nuclear factor- κ B phosphorylation and nuclear appearance in human endothelial cells, *Am. J. Clin. Nutr.* 77 (5) (2003) 1220–1228.
- [42] L.-M. Hung, J.-K. Chen, S.-S. Huang, R.-S. Lee, M.-J. Su, Cardioprotective effect of resveratrol, a natural antioxidant derived from grapes, *Cardiovasc. Res.* 47 (3) (2000) 549–555.
- [43] Z.-H. Ma, Q.-Y. Ma, L.-C. Wang, H.-C. Sha, S.-L. Wu, M. Zhang, Effect of resveratrol on peritoneal macrophages in rats with severe acute pancreatitis, *Inflamm. Res.* 54 (12) (2005) 522–527.
- [44] S.H. Tsai, S.Y. Lin-Shiau, J.K. Lin, Suppression of nitric oxide synthase and the down-regulation of the activation of NF κ B in macrophages by resveratrol, *Br. J. Pharmacol.* 126 (3) (1999) 673–680.
- [45] Y. Sugiyama, S. Kato, S. Mitsufoji, T. Okanoue, K. Takeuchi, Pathogenic role of endothelial nitric oxide synthase (eNOS/NOS-III) in cerulein-induced rat acute pancreatitis, *Dig. Dis. Sci.* 51 (8) (2006) 1396–1403.
- [46] K. Li, J. Yao, L. Shi, N. Sawada, Y. Chi, Q. Yan, H. Matsue, M. Kitamura, M. Takeda, Reciprocal regulation between proinflammatory cytokine-induced inducible NO synthase (iNOS) and connexin43 in bladder smooth muscle cells, *J. Biol. Chem.* 286 (48) (2011) 41552–41562.
- [47] S.R. Mettu, J.D. Wig, M. Khullar, G. Singh, R. Gupta, Efficacy of serum nitric oxide level estimation in assessing the severity of necrotizing pancreatitis, *Pancreatology* 3 (6) (2003) 506–514.
- [48] Y. Hoshino, H. Sonoda, R. Nishimura, K. Mori, K. Ishibashi, M. Ikeda, Involvement of the NADPH oxidase 2 pathway in renal oxidative stress in Aqp11 $^{-/-}$ mice, *Biochem. Biophys. Res. Commun.* 17 (2019) 169–176.
- [49] W. Deng, A. Abliz, S. Xu, R. Sun, W. Guo, Q. Shi, J. Yu, W. Wang, Severity of pancreatitis-associated intestinal mucosal barrier injury is reduced following treatment with the NADPH oxidase inhibitor apocynin, *Mol. Med. Rep.* 14 (4) (2016) 3525–3534.
- [50] Y. Wen, R. Liu, N. Lin, H. Luo, J. Tang, Q. Huang, H. Sun, L. Tang, NADPH oxidase hyperactivity contributes to cardiac dysfunction and apoptosis in rats with severe experimental pancreatitis through ROS-mediated MAPK signaling pathway, *Oxid. Med. Cell. Longev.* 2019 (2019).
- [51] N. Kim, J.M. Park, S.H. Lee, B.H. Kim, J.H. Son, J.K. Ryu, Y.-T. Kim, W. Lee, Effect of combinatory treatment with resveratrol and guggulsterone on mild acute pancreatitis in mice, *Pancreas* 46 (3) (2017) 366–371.
- [52] A. Szabolcs, I.S. Varga, C. Varga, A. Berkó, J. Kaszaki, T. Letoha, L. Tiszlavicz, R. Sári, J. Lonovics, T. Takács, Beneficial effect of resveratrol on cholecystokinin-induced experimental pancreatitis, *Eur. J. Pharmacol.* 532 (1–2) (2006) 187–193.
- [53] X.-P. Zhang, Z.-J. Li, J. Zhang, Inflammatory mediators and microcirculatory disturbance in acute pancreatitis, *Hepatobiliary Pancreat. Dis. Int.* 8 (4) (2009) 351–357.
- [54] M.P. Fuggetta, V. Bordignon, A. Cottarelli, B. Macchi, C. Frezza, P. Cordiali-Fei, F. Ensoli, S. Ciafrè, F. Marino-Merlo, A. Mastino, Downregulation of proinflammatory cytokines in HTLV-1-infected t cells by resveratrol, *J. Exp. Clin. Cancer Res.* 35 (1) (2016) 118.
- [55] N. Silswal, A.K. Singh, B. Aruna, S. Mukhopadhyay, S. Ghosh, N.Z. Ehtesham, Human resistin stimulates the pro-inflammatory cytokines TNF- α and IL-12 in macrophages by NF- κ B-dependent pathway, *Biochem. Biophys. Res. Commun.* 334 (4) (2005) 1092–1101.
- [56] T. Kasama, Y. Miwa, T. Isozaki, T. Odoi, M. Adachi, S.L. Kunkel, Neutrophil-derived cytokines: potential therapeutic targets in inflammation, *Curr. Drug Targets-Inflammation Allergy* 4 (3) (2005) 273–279.
- [57] J.W. Larrick, S.C. Wright, Cytotoxic mechanism of tumor necrosis factor- α , *FASEB J.* 4 (14) (1990) 3215–3223.
- [58] P.A. Baeuerle, T. Henkel, Function and activation of NF- κ B in the immune system, *Annu. Rev. Immunol.* 12 (1) (1994) 141–179.
- [59] A.S. Baldwin, Series introduction: the transcription factor NF- κ B and human disease, *J. Clin. Invest.* 107 (1) (2001) 3–6.
- [60] M. Jafarian, S.-H. Mozhgani, E. Patrad, H. Vaziri, S.A. Rezaee, M.M. Akbarin, M. Norouzi, Evaluation of iNOS, ICAM-1, and VCAM-1 gene expression: a study of adult T cell leukemia malignancy associated with HTLV-1, *Arch. Virol.* 162 (4) (2017) 1009–1015.
- [61] J.-K. Min, Y.-M. Kim, S.W. Kim, M.-C. Kwon, Y.-Y. Kong, I.K. Hwang, M.H. Won, J. Rho, Y.-G. Kwon, TNF-related activation-induced cytokine enhances leukocyte adhesiveness: induction of ICAM-1 and VCAM-1 via TNF receptor-associated factor and protein kinase C-dependent NF- κ B activation in endothelial cells, *J. Immunol.* 175 (1) (2005) 531–540.
- [62] R.J. Playford, A. Belo, R. Poulsom, A.J. Fitzgerald, K. Harris, I. Pawluczuk, J. Ryon, T. Darby, M. Nilsen-Hamilton, S. Ghosh, Effects of mouse and human lipocalin homologues 24p3/lcn2 and neutrophil gelatinase-associated lipocalin on gastrointestinal mucosal integrity and repair, *Gastroenterology* 131 (3) (2006) 809–817.
- [63] Y. Seo, J. Park, W. Choi, D. Ju Son, Y. Sung Kim, M.-K. Kim, B.-E. Yoon, J. Pyee, J. Tae Hong, Y.-M. Go, Antiatherogenic effect of resveratrol attributed to decreased expression of ICAM-1 (Intercellular adhesion Molecule-1) mechanistic link from focal adhesion to monocyte adhesion, *Arterioscler. Thromb. Vasc. Biol.* 39 (4) (2019) 675–684.
- [64] C.-W. Liu, H.-C. Sung, S.-R. Lin, C.-W. Wu, C.-W. Lee, I.-T. Lee, Y.-F. Yang, I.-S. Yu, S.-W. Lin, M.-H. Chiang, Resveratrol attenuates ICAM-1 expression and monocyte adhesiveness to TNF- α -treated endothelial cells: evidence for an anti-inflammatory cascade mediated by the miR-221/222/AMPK/p38/NF- κ B pathway, *Sci. Rep.* 7 (2017) 44689.
- [65] W. Pan, H. Yu, S. Huang, P. Zhu, Resveratrol protects against TNF- α -induced injury in human umbilical endothelial cells through promoting sirtuin-1-induced repression of NF- κ B and p38 MAPK, *PLoS One* 11 (1) (2016).
- [66] D. Tewari, S.F. Nabavi, S.M. Nabavi, A. Sureda, A.A. Farooqi, A.G. Atanasov, R. A. Vacca, G. Sethi, A. Bishayee, Targeting activator protein 1 signaling pathway by bioactive natural agents: possible therapeutic strategy for cancer prevention and intervention, *Pharmacol. Res.* 128 (2018) 366–375.
- [67] A. Gulcubuk, D. Haktanir, A. Cakiris, D. Ustek, O. Guzel, M. Erturk, F. Yildirim, I. Akyazi, H. Cicekci, M.H. Durak, The effects of resveratrol on tissue injury, oxidative damage, and pro-inflammatory cytokines in an experimental model of acute pancreatitis, *J. Physiol. Biochem.* 70 (2) (2014) 397–406.
- [68] X. Gao, Y.X. Xu, N. Janakiraman, R.A. Chapman, S.C. Gautam, Immunomodulatory activity of resveratrol: suppression of lymphocyte proliferation, development of cell-mediated cytotoxicity, and cytokine production, *Biochem. Pharmacol.* 62 (9) (2001) 1299–1308.
- [69] I. Gukovsky, C.N. Reyes, E.C. Vaquero, A.S. Gukovskaya, S.J. Pandol, Curcumin ameliorates ethanol and nonethanol experimental pancreatitis, *Am. J. Physiol. Gastrointestinal Liver Physiol.* 284 (1) (2003) G85–G95.
- [70] C.-M. Yang, Y.-W. Chen, P.-L. Chi, C.-C. Lin, L.-D. Hsiao, Resveratrol inhibits BK-induced COX-2 transcription by suppressing acetylation of AP-1 and NF- κ B in human rheumatoid arthritis synovial fibroblasts, *Biochem. Pharmacol.* 132 (2017) 77–91.
- [71] D. Liu, G. Song, Z. Ma, X. Geng, Y. Dai, T. Yang, H. Meng, J. Gong, B. Zhou, Z. Song, Resveratrol improves the therapeutic efficacy of bone marrow-derived mesenchymal stem cells in rats with severe acute pancreatitis, *Int. Immunopharmacol.* 80 (2020), 106128.
- [72] J.-y. Li, W.-q. Huang, R.-h. Tu, G.-q. Zhong, B.-b. Luo, Y. He, Resveratrol rescues hyperglycemia-induced endothelial dysfunction via activation of Akt, *Acta Pharmacol. Sin.* 38 (2) (2017) 182–191.
- [73] C.Y. Chang, D.-K. Choi, D.K. Lee, Y.J. Hong, E.J. Park, Resveratrol confers protection against rotenone-induced neurotoxicity by modulating myeloperoxidase levels in glial cells, *PLoS One* 8 (4) (2013).
- [74] M. Palacz-Wrobel, P. Borkowska, M. Paul-Samojedny, M. Kowalczyk, A. Fila-Danilow, R. Suchanek-Raif, J. Kowalski, Effect of apigenin, kaempferol and resveratrol on the gene expression and protein secretion of tumor necrosis factor

- alpha (TNF- α) and interleukin-10 (IL-10) in RAW-264.7 macrophages, *Biomed. Pharmacother.* 93 (2017) 1205–1212.
- [75] A.M. Wolf, D. Wolf, H. Rumpold, B. Enrich, H. Tilg, Adiponectin induces the anti-inflammatory cytokines IL-10 and IL-1RA in human leukocytes, *Biochem. Biophys. Res. Commun.* 323 (2) (2004) 630–635.
- [76] A. Piironen, Severe acute pancreatitis: contrast-enhanced CT and MRI features, *Abdom. Radiol.* 26 (3) (2001) 225.
- [77] K. Bosscha, P. Hulstaert, A. Hennipman, M. Visser, H. Gooszen, T.M. van Vroonhoven, C. vd Werken, Fulminant acute pancreatitis and infected necrosis: results of open management of the abdomen and “planned” reoperations, *J. Am. Coll. Surg.* 187 (3) (1998) 255–262.
- [78] Y. Meng, M. Zhang, J. Xu, X.-M. Liu, Q.-Y. Ma, Effect of resveratrol on microcirculation disorder and lung injury following severe acute pancreatitis in rats, *World J. Gastroenterol.* WJG 11 (3) (2005) 433.
- [79] R. Yang, J. Tenhunen, T.I. Tonnessen, HMGB1 and histones play a significant role in inducing systemic inflammation and multiple organ dysfunctions in severe acute pancreatitis, *Int. J. Inflam.* 2017 (2017).
- [80] L. Bonjoch, V. Casas, M. Carrascal, D. Closa, Involvement of exosomes in lung inflammation associated with experimental acute pancreatitis, *J. Pathol.* 240 (2) (2016) 235–245.
- [81] Z.S. Lin, C.F. Ku, Y.F. Guan, H.T. Xiao, X.K. Shi, H.Q. Wang, Z.X. Bian, S. W. Tsang, H.J. Zhang, Dihydro-resveratrol ameliorates lung injury in rats with cerulein-induced acute pancreatitis, *Phytother. Res.* 30 (4) (2016) 663–670.
- [82] A.S. Elder, G.T. Saccone, A.D. Bersten, D.-L. Dixon, Evaluation of lung injury and respiratory mechanics in a rat model of acute pancreatitis complicated with endotoxin, *Pancreatology* 12 (3) (2012) 240–247.
- [83] H. Wang, S. Wang, A. Tang, H. Gong, P. Ma, L. Chen, Combined effects of sivelestat and resveratrol on severe acute pancreatitis-associated lung injury in rats, *Exp. Lung Res.* 40 (6) (2014) 288–297.
- [84] Y.-Q. Zhang, Y.-J. Liu, Y.-F. Mao, W.-W. Dong, X.-Y. Zhu, L. Jiang, Resveratrol ameliorates lipopolysaccharide-induced epithelial mesenchymal transition and pulmonary fibrosis through suppression of oxidative stress and transforming growth factor- β 1 signaling, *Clin. Nutr.* 34 (4) (2015) 752–760.
- [85] C. Frey, H. Zhou, D. Harvey, R.H. White, Co-morbidity is a strong predictor of early death and multi-organ system failure among patients with acute pancreatitis, *J. Gastrointest. Surg.* 11 (6) (2007) 733–742.
- [86] G. Sun, F. Liu, C. Xiu, High thoracic sympathetic block improves coronary microcirculation disturbance in rats with chronic heart failure, *Microvasc. Res.* 122 (2019) 94–100.
- [87] C. Cuthbertson, C. Christophi, Disturbances of the microcirculation in acute pancreatitis, *Br. J. Surg.* 93 (5) (2006) 518–530.
- [88] H. Sha, Q. Ma, R.K. Jha, Z. Wu, Z. Qingyuan, Z. Wang, Z. Ma, X. Luo, C. Liu, Resveratrol suppresses microcirculatory disturbance in a rat model of severe acute pancreatitis, *Cell Biochem. Biophys.* 67 (3) (2013) 1059–1065.
- [89] R.K. Jha, Q. Ma, H. Sha, M. Palikhe, Protective effect of resveratrol in severe acute pancreatitis-induced brain injury, *Pancreas* 38 (8) (2009) 947–953.
- [90] Z. Li, Q. Ma, Y. Luo, Effect of resveratrol-induced FasL up-regulation on the apoptosis of pancreatic acinar cells in rats with severe acute pancreatitis, *Nan fang yi ke da xue xue bao* = J. Southern Med. Univ. 29 (3) (2009) 454–457.
- [91] H. Sha, Q. Ma, R.K. Jha, Z. Wang, Resveratrol ameliorates lung injury via inhibition of apoptosis in rats with severe acute pancreatitis, *Exp. Lung Res.* 35 (4) (2009) 344–358.
- [92] S.P. Ip, S.W. Tsang, T.P. Wong, C.T. Che, P.S. Leung, Saralasin, a nonspecific angiotensin II receptor antagonist, attenuates oxidative stress and tissue injury in cerulein-induced acute pancreatitis, *Pancreas* 26 (3) (2003) 224–229.
- [93] X.-P. Zhang, J. Zhang, M.-L. Ma, Y. Cai, R.-J. Xu, Q. Xie, X.-G. Jiang, Q. Ye, Pathological changes at early stage of multiple organ injury in a rat model of severe acute pancreatitis, *Hepatobiliary Pancreatic Dis. Int. HBPD INT* 9 (1) (2010) 83–87.
- [94] J. Jaworek, B. Jachimczak, R. Tomaszewska, P. Konturek, W. Pawlik, R. Sendur, E. Hahn, J. Stachura, S. Konturek, Protective action of lipopolysaccharides in rat cerulein-induced pancreatitis: role of nitric oxide, *Digestion* 62 (1) (2000) 1–13.
- [95] S. Rahman, B. Ammorio, M. Larvin, M. McMahon, Increased nitric oxide excretion in patients with severe acute pancreatitis: evidence of an endotoxin mediated inflammatory response? *Gut* 52 (2) (2003) 270–274.
- [96] R. Al-Mufti, R. Williamson, R. Mathie, Increased nitric oxide activity in a rat model of acute pancreatitis, *Gut* 43 (4) (1998) 564–570.
- [97] A. Satoh, T. Shimosegawa, K. Kimura, S. Moriizumi, A. Masamune, M. Koizumi, T. Toyota, Nitric oxide is overproduced by peritoneal macrophages in rat taurocholate pancreatitis: the mechanism of inducible nitric oxide synthase expression, *Pancreas* 17 (4) (1998) 402–411.
- [98] D. Qian, G. Song, Z. Ma, G. Wang, L. Jin, M. Hu, Z. Song, X. Wang, MicroRNA-9 modified bone marrow-derived mesenchymal stem cells (BMSCs) repair severe acute pancreatitis (SAP) via inducing angiogenesis in rats, *Stem Cell Res. Ther.* 9 (1) (2018) 1–15.
- [99] Y.S. Kim, J.H. Chang, T.H. Kim, C.W. Kim, J.K. Kim, S.W. Han, Prolonged hyperamylasemia in patients with acute pancreatitis is associated with recurrence of acute pancreatitis, *Medicine* 99 (3) (2020), e18861.
- [100] I. Borovickova, N.R. Bhatt, G.P. Boran, P.F. Ridgway, Persistent chronic hyperamylasemia: clinical interpretation and diagnostic approach, *JOP* 17 (17) (2016) 349–358.
- [101] G. Lippi, M. Valentino, G. Cervellini, Laboratory diagnosis of acute pancreatitis: in search of the Holy Grail, *Crit. Rev. Clin. Lab. Sci.* 49 (1) (2012) 18–31.
- [102] P.A. Banks, M.L. Freeman, P.P.C.o.t.A.C.o. Gastroenterology, Practice guidelines in acute pancreatitis, *Am. J. Gastroenterol.* 101 (10) (2006) 2379–2400.
- [103] M. Isogai, A. Yamaguchi, A. Hori, Y. Kaneoka, LDH to AST ratio in biliary pancreatitis—a possible indicator of pancreatic necrosis: preliminary results, *Am. J. Gastroenterol.* 93 (3) (1998).
- [104] B.E. Muhs, S. Patel, H. Yee, S. Marcus, P. Shamamian, Inhibition of matrix metalloproteinases reduces local and distant organ injury following experimental acute pancreatitis, *J. Surg. Res.* 109 (2) (2003) 110–117.
- [105] N. Wang, F. Zhang, L. Yang, J. Zou, H. Wang, K. Liu, M. Liu, H. Zhang, X. Xiao, K. Wang, Resveratrol protects against L-arginine-induced acute necrotizing pancreatitis in mice by enhancing SIRT1-mediated deacetylation of p53 and heat shock factor 1, *Int. J. Mol. Med.* 40 (2) (2017) 427–437.
- [106] M. Bhatia, F.L. Wong, Y. Cao, H.Y. Lau, J. Huang, P. Puneet, L. Chevali, Pathophysiology of acute pancreatitis, *Pancreatology* 5 (2–3) (2005) 132–144.
- [107] K.M. McAndrews, V.S. LeBleu, R. Kalluri, SIRT1 regulates lysosome function and exosome secretion, *Dev. Cell* 49 (3) (2019) 302–303.
- [108] M. Bhatia, M. Brady, S. Shokuchi, S. Christmas, J.P. Neoptolemos, J. Slavin, Inflammatory mediators in acute pancreatitis, *J. Pathol.* 190 (2) (2000) 117–125.
- [109] K.M. McAndrews, V.S. LeBleu, R. Kalluri, SIRT1 regulates lysosome function and exosome secretion, *Dev. Cell* 49 (3) (2019) 302–303.
- [110] L. Luu, F. Dai, K. Prentice, X. Huang, A. Hardy, J.B. Hansen, Y. Liu, J. Joseph, M. Wheeler, The loss of Sirt1 in mouse pancreatic beta cells impairs insulin secretion by disrupting glucose sensing, *Diabetologia* 56 (9) (2013) 2010–2020.
- [111] M. Sulaiman, M.J. Matta, N. Sunderesan, M.P. Gupta, M. Periasamy, M. Gupta, Resveratrol, an activator of SIRT1, upregulates sarcoplasmic calcium ATPase and improves cardiac function in diabetic cardiomyopathy, *Am. J. Physiol.-Heart Circulatory Physiol.* 298 (3) (2010) H833–H843.
- [112] L. Vetterli, P. Maechler, Resveratrol-activated SIRT1 in liver and pancreatic β -cells: a Janus head looking to the same direction of metabolic homeostasis, *Aging (Albany NY)* 3 (4) (2011) 444.
- [113] L. Wu, L. Zhou, Y. Lu, J. Zhang, F. Jian, Y. Liu, F. Li, W. Li, X. Wang, G. Li, Activation of SIRT1 protects pancreatic β -cells against palmitate-induced dysfunction, *Biochimica et Biophysica Acta (BBA)-Mol. Basis Dis.* 1822 (11) (2012) 1815–1825.
- [114] J.-H. Lee, M.-Y. Song, E.-K. Song, E.-K. Kim, W.S. Moon, M.-K. Han, J.-W. Park, K.-B. Kwon, B.-H. Park, Overexpression of SIRT1 protects pancreatic β -cells against cytokine toxicity by suppressing the nuclear factor- κ B signaling pathway, *Diabetes* 58 (2) (2009) 344–351.
- [115] G. Luo, L. Xiao, D. Wang, N. Wang, C. Luo, X. Yang, L. Hao, Resveratrol protects against ethanol-induced impairment of insulin secretion in INS-1 cells through SIRT1-UCP2 axis, *Toxicol. Vitro* 65 (2020), 104808.
- [116] Y. Nakamura, J.H. Do, J. Yuan, I.V. Odinokova, O. Mareninova, A.S. Gukovskaya, S.J. Pandol, Inflammatory cells regulate p53 and caspases in acute pancreatitis, *Am. J. Physiol.-Gastrointestinal Liver Physiol.* 298 (1) (2010) G92–G100.
- [117] L. Golkar, X.-Z. Ding, M.B. Ujiki, M.R. Salabat, D.L. Kelly, D. Scholtens, A. J. Fought, D.J. Bentrem, M.S. Talamonti, R.H. Bell, Resveratrol inhibits pancreatic cancer cell proliferation through transcriptional induction of macrophage inhibitory cytokine-1, *J. Surg. Res.* 138 (2) (2007) 163–169.
- [118] H. Zhang, L. Zhang, F. Yu, Y. Liu, Q. Liang, G. Deng, G. Chen, M. Liu, X. Xiao, HSF1 is a transcriptional activator of IL-10 gene expression in RAW264. 7 macrophages, *Inflammation* 35 (4) (2012) 1558–1566.
- [119] S. Chen, X. Zuo, M. Yang, H. Lu, N. Wang, K. Wang, Z. Tu, G. Chen, M. Liu, K. Liu, Severe multiple organ injury in HSF1 knockout mice induced by lipopolysaccharide is associated with an increase in neutrophil infiltration and surface expression of adhesion molecules, *J. Leukoc. Biol.* 92 (4) (2012) 851–857.
- [120] Z. Tong, B. Jiang, L. Zhang, Y. Liu, M. Gao, Y. Jiang, Y. Li, Q. Lu, Y. Yao, X. Xiao, HSF-1 is involved in attenuating the release of inflammatory cytokines induced by LPS through regulating autophagy, *Shock* 41 (5) (2014) 449–453.
- [121] A. Kumar, A.M. Rimando, A.S. Levenson, Resveratrol and pterostilbene as a microRNA-mediated chemopreventive and therapeutic strategy in prostate cancer, *Ann. N. Y. Acad. Sci.* 1403 (1) (2017) 15–26.
- [122] Y. Xin, H. Zhang, Z. Jia, X. Ding, Y. Sun, Q. Wang, T. Xu, Resveratrol improves uric acid-induced pancreatic β -cells injury and dysfunction through regulation of miR-126, *Biomed. Pharmacother.* 102 (2018) 1120–1126.
- [123] S.-t. Tang, F. Wang, M. Shao, Y. Wang, H.-q. Zhu, MicroRNA-126 suppresses inflammation in endothelial cells under hyperglycemic condition by targeting HMGB1, *Vascul. Pharmacol.* 88 (2017) 48–55.
- [124] J.P. Huddleson, Regulation of the KLF2 Transcription Factor Gene in Endothelial Cells by Fluid Shear Stress, University of Cincinnati, 2006.
- [125] S. Ali, R. Suresh, S. Banerjee, B. Bao, Z. Xu, J. Wilson, P.A. Philip, M. Apte, F. H. Sarkar, Contribution of microRNAs in understanding the pancreatic tumor microenvironment involving cancer associated stellate and fibroblast cells, *Am. J. Cancer Res.* 5 (3) (2015) 1251.
- [126] B. Yan, L. Cheng, Z. Jiang, K. Chen, C. Zhou, L. Sun, J. Cao, W. Qian, J. Li, T. Shan, Resveratrol inhibits ROS-promoted activation and glycolysis of pancreatic stellate cells via suppression of miR-21, *Oxid. Med. Cell. Longev.* 2018 (2018).
- [127] P. Liu, H. Liang, Q. Xia, P. Li, H. Kong, P. Lei, S. Wang, Z. Tu, Resveratrol induces apoptosis of pancreatic cancers cells by inhibiting miR-21 regulation of BCL-2 expression, *Clin. Transl. Oncol.* 15 (9) (2013) 741–746.
- [128] S. Feng, Q. Wei, Q. Hu, X. Huang, X. Zhou, G. Luo, M. Deng, M. Lü, Research progress on the relationship between acute pancreatitis and calcium overload in acinar cells, *Dig. Dis. Sci.* 64 (1) (2019) 25–38.
- [129] L. Wen, R. Mukherjee, W. Huang, R. Sutton, Calcium Signaling, Mitochondria and Acute Pancreatitis: Avenues for Therapy, *Pancreapedia: the Exocrine Pancreas Knowledge Base*, 2016.

- [130] G.L. Smith, D.A. Eisner, Calcium buffering in the heart in health and disease, *Circulation* 139 (20) (2019) 2358–2371.
- [131] T. Peng, X. Peng, M. Huang, J. Cui, Y. Zhang, H. Wu, C. Wang, Serum calcium as an indicator of persistent organ failure in acute pancreatitis, *Am. J. Emerg. Med.* 35 (7) (2017) 978–982.
- [132] L. Wang, Q. Ma, X. Chen, H. Sha, Z. Ma, Effects of resveratrol on calcium regulation in rats with severe acute pancreatitis, *Eur. J. Pharmacol.* 580 (1-2) (2008) 271–276.